



Memorial Regional Medical Center
 Rappahannock General Hospital
 Reynolds Crossing Imaging Center
 Richmond Community Hospital
 St. Francis Medical Center
 St. Mary's Hospital
 Midlothian Imaging
 Imaging Center at Innsbrook
 Westchester Imaging Center

INSTRUCTIONS FOR PATIENTS ON METFORMIN WHO RECEIVE IODINATED IV CONTRAST MEDIA

DRUGS CONTAINING METFORMIN INCLUDE:

ACTOPLUS MET	JENTADUETO
ACTOPLUS MET XR	JENTADUETO XR
AVANDAMET	KAZANO
FORTAMET	KOMBIGLYZE XR
GLUCOPHAGE	METAGLIP
GLUCOPHAGE XR	PRANDIMET
GLUCOVANCE	RIOMET
GLUMETZA	SYNJARDY
INVOKAMET	SYNJARDY XR
INCOKAMET XR	
JANUMET	
JANUMET XR	

- You should not take your metformin (e.g. Glucophage®) or any other metformin containing oral antidiabetic agents (*See above Chart*), **for 48 hours** following today's IV Contrast administration.
- You should check with your physician before restarting the metformin.
- Your doctor may want to draw your blood to check your kidney function before you resume taking your metformin.
- In addition, we advise that you drink plenty of fluids to adequately hydrate yourself over the next 24 hours.
- Any questions should be directed to your primary care physician.

Patient Signature: _____

Study performed: _____

Technologist: _____

Date: _____